

EFFECTIVE SHARED CARE AGREEMENT OF MYCOPHENOLATE FOR THE TREATMENT OF AUTOIMMUNE RHEUMATIC DISEASES

(use is off-label for this indication – but included in national BSR/NICE/Cochrane reviews and guidelines)

RESPONSIBILITIES and ROLES

Specialist clinician responsibilities	
1	Discuss the benefits and side effects of treatment with the patient.
2	Check for possible drug interactions with mycophenolate and patients regular medication. Avoid prescribing interacting drugs.
3	Assess likelihood of compliance.
4	Ensure patient is given sufficient information about their treatment.
5	Perform all baseline tests (including FBC, U&Es and LFTs).
6	Initiate and stabilise treatment with mycophenolate. Stabilisation will usually take 4 weeks.
7	Supply medication until care is transferred to GP.
8	Ask the GP whether he or she is willing to participate in shared care and explain the intention to share care with patient/carer and obtain consent.
9	Monitors appropriately as stated on the monitoring requirement.
10	Inform GP of the dose to be prescribed, any changes in dose, when to stop treatment and when to refer the patient back to specialist clinician.
11	Ensure GP has access to blood results for information.
12	Monitor for side effects and report adverse events to the MHRA and GP where appropriate.
13	Inform GP if patient does not attend specialist clinician appointments and action to be taken.
14	Have a mechanism in place to receive rapid referral of a patient from the GP in event of deteriorating clinical condition.
15	Ensure that clear backup arrangements exist for GPs to obtain advice and support.
General Practitioner responsibilities	
1	Reply to the request for shared care as soon as possible.
2	Prescribe mycophenolate at the dose recommended once patient is established on treatment.
3	Ensure compatibility with other concomitant medication.
4	Adjust the dose as advised by the specialist clinician.
5	Contact the specialist clinician if you suspect the patient is not complying with their medication.
6	Check for possible drug interaction when prescribing new medication and avoid prescribing interacting drugs.
7	Stop treatment on the advice of the specialist clinician or immediately if an urgent need to stop treatment arises.
8	Refer the patient to the specialist clinician if his/her condition deteriorates.
9	Report any suspected adverse events to specialist team and any severe adverse events to MHRA.
Patient's / Carer's role	
1	Take mycophenolate as recommended by specialist clinician.
2	Report to the specialist clinician or GP if he / she does not have a clear understanding of the treatment.
3	Request repeat prescriptions from the GP at least 5 days before the next supply is needed.
4	Attend scheduled appointments with specialist clinician, GP and for monitoring.
5	Share any concerns in relation to treatment with GP or specialist clinician.
6	Inform specialist clinician or GP of any other medication being taken, including over-the-counter products
7	Inform specialist clinician or GP if you feel you are having problems taking your medication or have stopped taking it.
8	Report any adverse effects to the specialist clinician or GP.

SUPPORTING INFORMATION EFFECTIVE SHARED CARE AGREEMENT

Information on therapeutic indication, dosage, method of administration, side effects and management considerations in special populations can be found in the Summary of Product Characteristics for mycophenolate available from www.medicines.org.uk

Monitoring requirements

Monitoring Interval	Full blood count	LFTs	U&Es	Albumin	GP or Specialist clinician
Baseline	✓	✓	✓	✓	Specialist clinician
Every 2 weeks during dose titration and for 6 weeks at stable dose	✓	✓	✓	✓	Specialist clinician
Every month for 3 months	✓	✓	✓	✓	Specialist clinician
Every 3 months	✓	✓	✓	✓	Specialist clinician

Additional information which cannot be found in the SPC

Reference

[BSR and BHPR guideline for the prescription and monitoring of non- biologic disease-modifying anti-rheumatic drugs 2017 Rheumatology 56:865-868](#)

<https://www.nice.org.uk/advice/esuom36/chapter/Key-points-from-the-evidence>

<https://www.nice.org.uk/advice/esuom32/chapter/Key-points-from-the-evidence>

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD002922.pub3/abstract>

BACK-UP ADVICE AND SUPPORT

Contact details	Telephone No.
Helpline:	01782 673687
For urgent advice ask for Rheumatology consultant or SpR on-call	01782 715444
Pharmacy (MPFT):	01782 673767